

Geometry of genus 9 Fano 4-fold

Han Frédéric

Institut de Mathématiques de Jussieu,
 Université Paris 7 - Case Postale 7012,
 2 Place Jussieu
 75251 Paris Cedex 05 France
 email: han@math.jussieu.fr

Introduction:

Let W be a 6 dimensional vector space over the complex number, equipped with a non degenerate symplectic form ω . Let G_ω be the grassmannian of ω -isotropic 3-dimensional vector subspaces of W . Considering the Plucker embedding, the intersection of G_ω with a generic codimension 2 linear subspace is the Mukai model of a smooth Fano manifold of dimension 4, genus 9 index 2 and picard number 1. We will also note by P_ω the 13 dimensional projective space spanned by G_ω under it's plucker embedding.

NOTATIONS:

In all the paper, B will be a general double hyperplane section of G_ω . For a hyperplane H of P_ω , we define $\bar{H} = H \cap G_\omega$, and for any $u \in G_\omega$, the corresponding plane of $\mathbb{P}(W)$ will be noted π_u .

Abstract

On a genus 9 Fano variety, Mukai's construction gives a natural rank 3 vector bundle, but curiously in dimension 4, another phenomena appears. In the first part of this article, we will explain how to construct on a Fano 4-fold of genus 9 (named B), a canonical set of four stable vector bundles of rank 2, and prove that they are rigid. Those bundles was already known to A. Iliev and K. Ranestad in [I-R] and the results of this section could be viewed as particular cases of the work of A. Kuznetsov (Cf [K]). We'd like here to show their consequences in the geometry of the 4-fold, and study Zak duality in this case.

Indeed, this "four-ality" (Cf [M]) is also present in the geometry of lines included in B , and also in the Chow ring of B . In section 2 we show that the variety of lines in B , is an hyperplane section of $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$. This description is explicit and could also be interesting in terms of Freudenthal geometries. Then in section 3, we compute the Chow ring of B which appears to have a rich structure in codimension 2.

The 4 bundles constructed can embed B in a Grassmannian $G(2, 6)$, and the link with the order one congruence discovered by E. Mezzeti and P de Poi in [M-dP] will be done in section 4. In particular we will prove that the generic fano variety of genus 9 and dimension 4 can be obtained by their construction, and explain the choices involved. We will also describe in this part the normalization of the non quadratically normal variety they constructed, and also its variety of plane cubics.

Acknowledgements:

I'd like to thanks L. Gruson for his constant interest on this work, and also F. Zak and C. Peskine for fruitful discussions. I'd like also to thanks K. Ranestad and A. Kuznetsov for pointing and explaining their works.

1 Construction of rank 2 vector bundles on B

This part is devoted to the construction of a canonical set of 4 stable and rigid rank 2 vector bundles on B . Many results are done in a universal way in [K], but we detail this description to use it in next sections. Let's first recall some classical geometric properties of G_ω . (Cf [I]) The union of the tangent spaces to G_ω is a quartic hypersurface of P_ω , so a general line of P_ω has naturally 4 marked points. Dually, as the variety B is given by a pencil L of hyperplane sections of G_ω , there are in this pencil, 4 hyperplanes $H_1, \dots, H_4$ tangent to G_ω . Denoting by u_i the contact point of H_i with G_ω , we will first construct a rank 2 sheaf on $H_i \cap G_\omega$ with singular locus u_i , and it's restriction to B will be the vector bundle.

1.1 Data associated to a tangent hyperplane section

Let $u \in G_\omega$, and H be a general hyperplane tangent to G_ω at u . For any v in G_ω , denote by π_v the corresponding projective subspace of $\mathbb{P}(W)$, and consider the hyperplane section of G_ω :

$$\bar{H}_u = \{v \in G_\omega, \pi_v \cap \pi_u \neq \emptyset\}$$

It's proved in [I] the following:

Lemma 1.1 *There is a conic C in π_u such that $v \in H \cap \bar{H}_u \iff \pi_v \cap C \neq \emptyset$. For H general containing the tangent space of G_ω at u , C is smooth. Furthermore, $H \cap \bar{H}_u$ contains the tangent cone $T_u G_\omega \cap G_\omega = \{v \in G_\omega \mid \dim \pi_v \cap \pi_u \geq 1\}$ which is embedded in P_ω as a cone over a veronese surface.*

Let Z be the following incidence:

$$Z_H = \{(p, v) \in C \times (H \cap G_\omega) \mid p \in \pi_v\}$$

Identifying C with $\mathbb{P}_1$, we denote by q_1 and q_2 the projections from $C \times G_\omega$ to $\mathbb{P}_1$ and to G_ω , and by L the SL_2 -representation $H^0 \mathcal{O}_{\mathbb{P}_1}(1)$. Restricting the surjection $L \otimes \mathcal{O}_{\bar{H}} \rightarrow q_{2*} \mathcal{O}_{Z_H}(1, 0)$ to the hyperplane section $\bar{H}$, we obtain:

Proposition 1.2 *The sheaf $\mathcal{E}$ defined by the following exact sequence:*

$$0 \rightarrow \mathcal{E} \rightarrow L \otimes \mathcal{O}_{\bar{H}} \rightarrow q_{2*} \mathcal{O}_{Z_H}(1, 0) \rightarrow 0$$

is reflexive of rank 2, $c_1(\mathcal{E}) = -1$ and is locally free outside u .

Proof: From lemma 1.1, the support of $q_{2*}\mathcal{O}_{Z_H}(1,0)$ is $H \cap \bar{H}_u$ so it is an hyperplane section of $\bar{H}$, hence $\mathcal{E}$ has rank 2 and $c_1(\mathcal{E}) = -1$. Furthermore, for v in $q_2(Z_H)$, the fiber of the restriction of q_2 to Z_H : $q_{2|Z_H}^{-1}(v)$ has length 1 if v is not in $\bar{H}_u$, it has length 2 if $v \in \bar{H}_u - \{u\}$ and $q_{2|Z_H}^{-1}(u)$ is the curve C . As Z_H and $\bar{H} - u$ are smooth, the sheaf $q_{2*}\mathcal{O}_{Z_H}(1,0)$ has projective dimension 1 outside u , hence $\mathcal{E}$ is locally free outside u . $\square$

Denote by $S_i L$ the SL_2 -representation $H^0(\mathcal{O}_{\mathbb{P}_1}(i))$ and by K and Q the tautological bundles on G_ω , such that the following sequence is exact.

$$0 \rightarrow K \rightarrow W \otimes \mathcal{O}_{G_\omega} \rightarrow Q \rightarrow 0$$

Proposition 1.3 *For $i > 0$ we have $R^i q_{2*}\mathcal{O}_{Z_H}(1,0) = 0$, and the resolution of $q_{2*}\mathcal{O}_{Z_H}(1,0)$ in G_ω is given by the following exact sequence:*

$$0 \rightarrow S_3 L \otimes \mathcal{O}_{G_\omega}(-1) \rightarrow L \otimes \bigwedge^2 Q^\vee \rightarrow L \otimes \mathcal{O}_{G_\omega} \rightarrow q_{2*}\mathcal{O}_{Z_H}(1,0) \rightarrow 0 \quad (1)$$

Proof: We consider the injection from $q_1^*(\mathcal{O}_{\mathbb{P}_1}(-2))$ to $W \otimes \mathcal{O}_{\mathbb{P}_1 \times G_\omega}$ given by the conic C . So the incidence Z_H is the locus where the map from $q_1^*(\mathcal{O}_{\mathbb{P}_1}(-2)) \oplus q_2^* K$ to $W \otimes \mathcal{O}_{\mathbb{P}_1 \times G_\omega}$ is not injective, hence Z_H is obtained in $\mathbb{P}_1 \times G_\omega$ as the zero locus of a section of the bundle $\mathcal{O}_{\mathbb{P}_1}(2) \boxtimes Q$.

Let $\mathcal{K}^\cdot$ be the Koszul complex $\bigwedge^i(\mathcal{O}_{\mathbb{P}_1}(-2) \boxtimes Q^\vee)$ of this section. We obtain the proposition 1.3 from the Leray spectral sequence applied to $\mathcal{K}^\cdot$ twisted by $\mathcal{O}_{\mathbb{P}_1 \times G_\omega}(1,0)$. $\square$

Furthermore, we deduce from Bott's theorem on G_ω the following:

Corollary 1.4 *We have the following equality $L = H^0(\mathcal{O}_{Z_H}(1,0) = H^0(q_{2*}\mathcal{O}_{Z_H}(1,0))$, and for $i > 0$, all the groups $H^i(\mathcal{O}_{Z_H}(1,0))$ and $H^i(q_{2*}\mathcal{O}_{Z_H}(1,0))$ are zero. For $i \geq 0$ all the groups $H^i(q_{2*}\mathcal{O}_{Z_H}(1,-1))$ and $H^i(q_{2*}\mathcal{O}_{Z_H}(1,-1))$ are zero.*

Proof: We will prove that on the isotropic Grassmannian G_ω , the bundles $\bigwedge^i Q^\vee$ and $(\bigwedge^i Q^\vee)(-1)$ are acyclic for $i \in \{1, 2, 3\}$. Indeed, with the notations of [W] 4.3.3 and 4.3.4, they correspond to the partitions $(0, 0, -1)$, $(0, -1, -1)$, $(-1, -1, -1)$, $(-1, -1, -2)$, $(-1, -2, -2)$, $(-2, -2, -2)$. Now recall that the half sum of positive roots is $\rho = (3, 2, 1)$, so $\alpha + \rho$ either contains a 0 or is $(2, 1, -1)$. So in all cases $\alpha + \rho$ is invariant by a signed permutation, and the sheaves are acyclic. The corollary is now a direct consequence of this acyclicity. $\square$

Corollary 1.5 *The sheaves $\mathcal{E}$ and $\mathcal{E}(-1)$ are acyclic. The vector space $V = H^0(\mathcal{E}(1))$ has dimension 6 and $\forall i > 0, h^i(\mathcal{E}(1)) = 0$, and $\mathcal{E}(1)$ is generated by its global sections.*

Proof: The acyclicity of $\mathcal{E}$ and $\mathcal{E}(-1)$ is a direct consequence of the definition of $\mathcal{E}$ and of the previous corollary.

To obtain the second assertion, we restrict the sequence 1 to the hyperplane section $\bar{H}$, so we obtain the following monad¹:

$$0 \rightarrow S_3 L \otimes \mathcal{O}_{\bar{H}}(-1) \rightarrow L \otimes \bigwedge^2 Q_{\bar{H}}^\vee \rightarrow \mathcal{E} \rightarrow 0$$

whose cohomology is $\mathcal{T}or^1(q_{2*}(\mathcal{O}_{Z_H}(1, 0)), \mathcal{O}_{\bar{H}})$ which is equal to $q_{2*}(\mathcal{O}_{Z_H}(1, -1))$ because $Z_H \subset q_2^{-1}(H)$. Twisting this monad by $\mathcal{O}_{\bar{H}}(1)$ we obtain that $H^0(\mathcal{E}(1))$ is the quotient of $L \otimes W$ by $S_3 L \oplus L$ because $W = H^0(Q_{\bar{H}})$. Furthermore, the right part of the monad gives a surjection from $L \otimes Q_{\bar{H}}$ to $\mathcal{E}(1)$. But $L \otimes Q_{\bar{H}}$ is globally generated, so $\mathcal{E}(1)$ is also generated by its global sections.

The vanishing of $h^i(\mathcal{E}(1))$ for $i > 0$ is a corollary of the vanishing of $h^i(q_{2*}(\mathcal{O}_{Z_H}(1, 0)))$, $h^i(Q_{\bar{H}})$ and $h^i(\mathcal{O}_{\bar{H}})$ for $i > 0$. $\square$

We can remark that the two vector spaces V and W of dimension 6 have not the same role. More precisely, the conic C gives a marked subspace of W so that we have the following:

Remark 1.6 *The tangent hyperplane H gives canonically the SL_2 -equivariant sequences:*

$$0 \rightarrow S_2 L \rightarrow W \rightarrow S_2 L \rightarrow 0 \text{ and } 0 \rightarrow L \rightarrow V \rightarrow S_3 L \rightarrow 0$$

1.2 The 4 rank 2 vector bundles on B

The pencil of hyperplanes defining B contains the 4 tangent hyperplanes H_i , so we can apply the previous construction to construct a rank 2 sheaf $\mathcal{E}_i$ on each of the $\bar{H}_i$, and define by E_i the restriction of $\mathcal{E}_i$ to B . Because B is smooth, it doesn't contain the contact points u_i , so E_i is locally free on B .

Corollary 1.7 *All the cohomology groups of the vector bundles E_i vanish. In particular, the rank 2 vector bundles E_i are stable. The vector space $H^0(E_i(1))$ has dimension 6, and $\forall j > 0, h^j(E_i(1)) = 0$. The bundles $E_i(1)$ are generated by their global sections.*

Proof: It's a direct consequence of corollary 1.5, because B is a hyperplane section of $\bar{H}_i$. (Note that the stability condition for a E_i is equivalent to $h^0 E_i = 0$) $\square$

1.3 The restricted incidences

Now, for each of the 4 hyperplanes H_i containing B and tangent to G_ω at some point u_i , let C_i be the conic of the projective plane π_{u_i} constructed in 1.1. Consider the restriction of the incidences Z_{H_i} to B . In other words, let Z_i, Z'_i be:

$$Z_i = \{(p, v) \in C_i \times B \mid p \in \pi_v\}, \quad Z'_i = \{(p, v) \in Z_i \mid \dim(\pi_v \cap \pi_{u_i}) > 0\}$$

where q_1 and q_2 still denote the projections from $C_i \times G_\omega$ to C_i and G_ω .

Remark 1.8 *Let p be a fixed point of C_i . The scheme $Z_{i,p} = q_2(q_1^{-1}(p) \cap Z_i)$ is a 2 dimensional irreducible quadric in P_ω . The restriction of q_2 to Z'_i is a double cover of a veronese surface $V_i = q_2(Z'_i)$.*

¹A monad is complex exact at all terms different from the middle one.

Proof: In fact $\{b \in G_\omega | p \in \pi_v\}$ is a smooth quadric of dimension 3 (Cf [I]), so it doesn't contain planes. This scheme is included in H_i , so $Z_{i,p}$ is just an hyperplane section of this smooth quadric. It is also proved in [I] that $\{v \in G_\omega | \dim \pi_{u_i} \cap \pi_v > 0\}$ is a cône over a veronese surface of vertex u_i . As $u_i \notin B$, the surface V_i is the intersection of this cone with an hyperplane which doesn't contain the vertex u_i , so it's a veronese surface. $\square$

NOTATIONS:

We denote by σ_i the class of a point on C_i , and h_3 the class of a hyperplane in P_ω . (the plucker embedding of G_ω).

Proposition 1.9 *The incidence Z_i is a divisor of classe $2h$ in*

$$\Pi = Proj(\mathcal{O}_{\mathbb{P}_1}(2) \oplus S_2L \otimes \mathcal{O}_{\mathbb{P}_1})$$

Furthermore we have $h_3 \sim h + 2\sigma_i$, where $h \sim \mathcal{O}_\Pi(1)$ and σ_i is also the class of a point on the base of the fibration Π . The divisor Z'_i of Z_i is equivalent to $h - 2\sigma_i$.

Proof: Denotes by e_i the image of the map from $\mathcal{O}_{\mathbb{P}_1}(-2)$ to $W \otimes \mathcal{O}_{\mathbb{P}_1}$ associated to C_i . Choose an element ϕ' of $\bigwedge^3 W^\vee$ such that $\ker \phi'$ gives an hyperplane section of G_ω containing B and different from the $\bar{H}_i$. (i.e $\phi'(u_i) \neq 0$). We can remark that the incidence Z_i is given over $\mathbb{P}_1$ by the isotropic 2-dimensional subspaces l of $\frac{e_i^\perp}{e_i}$, such that $\phi'(e_i \otimes \wedge^2 l) = 0$, because the condition $\phi_i(e_i \otimes \wedge^2 l) = 0$ is already satisfied by the definition of C_i and lemma 1.1. (where ϕ_i denotes a trilinear form of kernel H_i)

The bundle $e_i^\perp$ is isomorphic to $S_2L \otimes \mathcal{O}_{\mathbb{P}_1} \oplus L \otimes \mathcal{O}_{\mathbb{P}_1}(-1)$ where the trivial factor S_2L correspond to the plane π_{u_i} . So² the bundle $\frac{e_i^\perp}{e_i}$ is isomorphic to $L(1) \oplus L(-1)$ where those factors are isotropic for the symplectic form induced by ω . We can take local basis s_0, s_1 and s_2, s_3 of each factors such that the form induced by ω is $p_{0,2} + p_{1,3}$ where $p_{i,j}$ denotes the Plucker coordinates associated to the s_i .

So the relative isotropic grassmannian $G_\omega(2, \frac{e_i^\perp}{e_i})$ is the intersection of $G(2, \frac{e_i^\perp}{e_i})$ with the subsheaf $\mathcal{O}_{\mathbb{P}_1}(2) \oplus \mathcal{O}_{\mathbb{P}_1}(-2) + S_2L \otimes \mathcal{O}_{\mathbb{P}_1}$ where the factor $\mathcal{O}_{\mathbb{P}_1}(2)$ still correspond to $s_0 \wedge s_1$.

Now we need to compute the kernel of the map $e_i \otimes \bigwedge^2(\frac{e_i^\perp}{e_i}) \xrightarrow{\phi'} \mathcal{O}_{\mathbb{P}_1}$. But the assumption $\phi'(u_i) \neq 0$ proves that it is $\mathcal{O}_{\mathbb{P}_1}(-4) \oplus L \otimes L \otimes \mathcal{O}_{\mathbb{P}_1}$.

So we have an exact sequence:

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_1}(-2) \oplus S_2L \otimes \mathcal{O}_{\mathbb{P}_1} \rightarrow \bigwedge^2(\frac{e_i^\perp}{e_i}) \xrightarrow{(\omega)_{\phi'}} \mathcal{O}_{\mathbb{P}_1} \oplus e_i^\vee \rightarrow 0$$

and Z_i is a divisor of class $2h$ in $Proj(\mathcal{O}_{\mathbb{P}_1}(2) \oplus S_2L \otimes \mathcal{O}_{\mathbb{P}_1})$. The relation $h_3 \sim h + 2\sigma_i$ is given by the map $e_i \otimes \bigwedge^2(\frac{e_i^\perp}{e_i}) \rightarrow \bigwedge^3 W$.

The divisor Z'_i of Z_i is locally given by the vanishing of the exterior product with $s_0 \wedge s_1$ so it equivalent to $h - 2\sigma_i$. $\square$

We will now study the relation between the conormal bundle of Z_i in $\mathbb{P}_1 \times B$ and the bundle E_i .

²As SL_2 -representation, we will identify L with its dual.

1.4 Deformations of E_i

Lemma 1.10 *We have the following exact sequence:*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}_1 \times B}(-\sigma_i - h_3) \rightarrow q_2^* E_i \rightarrow \mathcal{I}_{Z_i}(\sigma_i) \rightarrow q_2^*(R^1 q_{2*} \mathcal{I}_{Z_i})(-\sigma_i) \rightarrow 0$$

(where q_1 and q_2 denotes the projections from $\mathbb{P}_1 \times B$ to $\mathbb{P}_1$ and B)

Proof: From the resolution of the diagonal of $\mathbb{P}_1 \times \mathbb{P}_1$, we obtain the relative Beilinson's spectral sequence:

$$E_{a,b}^1 = (\bigwedge^{-a} \omega_{q_2}(\sigma_i)) \otimes R^b q_{2*}(\mathcal{I}_{Z_i}((1+a)\sigma_i)) \implies \mathcal{I}_{Z_i}(\sigma_i)$$

By the definition of E_i (Cf prop 1.2) we have $E_i = q_{2*}(\mathcal{I}_{Z_i}(\sigma_i))$. Furthermore, the projection $q_2(Z_i)$ is an hyperplane section of B , so $q_{2*} \mathcal{I}_{Z_i} = \mathcal{O}_B(-1)$. We can conclude, remarking that $R^1 q_{2*}(\mathcal{I}_{Z_i}(\sigma_i)) = 0$, because the restriction of $q_2: Z_i \xrightarrow{q_2|_{Z_i}} q_2(Z_i)$ has all its fibers of length at most 2. $\square$

NB: The support of $R^1 q_{2*} \mathcal{I}_{Z_i}$ is the natural scheme structure (Cf [G-P]) on the scheme of fibers of q_2 intersecting Z_i in length 2 or more. It is the veronese surface $V_i = q_2(Z'_i)$.

So the previous lemma can now be translated in the following:

Corollary 1.11 *The scheme $q_2^{-1}(V_i) \cup Z_i$ is in $\mathbb{P}_1 \times B$ the zero locus of a section of the bundle $q_2^* E_i(1, 1)$.*

This gives also a geometric description of the marked pencil of sections of $E_i(h_3)$ given by the natural inclusion $L \subset V$ found in remark 1.6. Indeed, if we fixe a point p on C_i , the restriction to $q_1^{-1}(p)$ of the section obtained in corollary 1.11 gives with the notations of lemma 1.8 the following:

Corollary 1.12 *For any point p on the conic C_i , the vector bundle $E_i(h_3)$ has a section vanishing on $Z_{i,p} \cup V_i$.*

We can now study the restriction of E_i to Z_i .

Proposition 1.13 *The restriction $E_{i|Z_i}$ of the vector bundle $q_2^* E_i$ to Z_i fits into the following exact sequence:*

$$0 \rightarrow \mathcal{O}_{Z_i}(h_3 - 3\sigma_i) \rightarrow q_2^* E_{i|Z_i}(h_3) \rightarrow \mathcal{O}_{Z_i}(3\sigma_i) \rightarrow 0$$

Proof: Fix a point p on C_i , and consider the corresponding section of $E_i(h_3)$ constructed in corollary 1.12. Its pull back gives a section of $q_2^* E_i(h_3)$ vanishing on $q_2^{-1}(Z_{i,p} \cup V_i)$, so its restriction to Z_i gives a section of $q_2^* E_{i|Z_i}(h_3 - \sigma_i - Z'_i)$. Now, using the computation of the class of Z'_i in Z_i made in proposition 1.9, namely that $\mathcal{O}_{Z_i}(Z'_i)$ is $\mathcal{O}_{Z_i}(h_3 - 4\sigma_i)$, it gives a section of $E_{i|Z_i}(3\sigma_i)$. We have to prove that it is a non vanishing section. To obtain this, we compute the second Chern's class of $E_{i|Z_i}(3\sigma_i)$. We will show that its image in the Chow ring of $\mathbb{P}_1 \times B$ is zero. Denote by a_i the second Chern class of E_i . From the lemma 1.11, we obtain the class of Z_i in $\mathbb{P}_1 \times B$: $[Z_i] = a_i + h_3 \cdot \sigma_i - [V_i]$. So we can compute $[Z_i].c_2(E_i(3\sigma_i))$. It is $(a_i + h_3 \cdot \sigma_i - [V_i]) \cdot (a_i - 3h_3 \sigma_i)$, but we will compute in proposition 3.7 the Chow ring of B , and this class vanish. $\square$

Corollary 1.14 *The vector bundles E_i are rigid, in other words we have $\text{Ext}^1(E_i, E_i) = 0$.*

Proof: From the corollary 1.11 we have an exact sequence on $\mathbb{P}_1 \times B$:

$$0 \rightarrow q_2^*E_i(-\sigma_i) \rightarrow q_2^*(E_i) \otimes q_2^*(E_i)(h_3) \rightarrow q_2^*E_i(h_3 + \sigma_i) \rightarrow (q_2^*E_i(h_3 + \sigma_i))|_{Z_i \cup q_2^{-1}(V_i)} \rightarrow 0$$

The bundle $q_2^*E_i(-\sigma_i)$ is acyclic, and the corollary 1.7 gives $H^0(q_2^*E_i(h_3 + \sigma_i)) = L \otimes V$ and $H^1(q_2^*E_i(h_3 + \sigma_i)) = 0$. The liaison exact sequence twisted by $q_2^*(E_i(h_3 + \sigma))$ is:

$$0 \rightarrow q_2^*E_i(h_3) \otimes \mathcal{O}_{q_2^{-1}(V_i)}(\sigma_i - Z'_i) \rightarrow q_2^*E_i(h_3 + \sigma_i)|_{Z_i \cup q_2^{-1}(V_i)} \rightarrow E_i|_{Z_i}(h_3 + \sigma_i) \rightarrow 0$$

As $\sigma_i - Z'_i$ have degree -1 along the fibers of $q_2 : q_2^{-1}(V_i) \rightarrow V_i$, the bundle $q_2^*E_i(h_3) \otimes \mathcal{O}_{q_2^{-1}(V_i)}(\sigma_i - Z'_i)$ is acyclic, so the cohomology of $q_2^*E_i(h_3 + \sigma_i)|_{Z_i \cup q_2^{-1}(V_i)}$ can be computed with its restriction to Z_i . The propositions 1.13 and 1.9 show that $H^0 E_i|_{Z_i}(h_3 + \sigma_i) = S_2 L \oplus S_2 L \oplus S_4 L$. In conclusion, we have the exact sequence:

$$0 \rightarrow \text{Hom}(E_i, E_i) \rightarrow L \otimes V \rightarrow S_2 L \oplus S_2 L \oplus S_4 L \rightarrow \text{Ext}^1(E_i, E_i) \rightarrow 0$$

By the corollary 1.7, the bundle E_i is stable, so it is simple, in other words we have $\text{Hom}(E_i, E_i) = \mathbb{C}$, and the above exact sequence gives $\text{Ext}^1(E_i, E_i) = 0$. $\square$

2 The variety of lines in B

Remark 2.1 *Let δ be an isotropic line of $\mathbb{P}(W)$. The set of isotropic planes of $\delta^\perp$ containing δ form a line in $G_\omega(3, 6)$, and all the line in $G_\omega(3, 6)$ are of this type for a unique element of $G_\omega(2, 6)$. In other words, the variety of lines in $G_\omega(3, 6)$ is naturally $G_\omega(2, 6)$.*

NOTATIONS:

A point of $G_\omega(2, 6)$ will be denoted by a minuscule letter, and the corresponding line in $G_\omega(3, 6)$ by the majuscule letter. The variety of lines included in B will be noted F_B . Denote by I the incidence point/line in B . In other words:

$$I = \text{Proj}\left(\left(\frac{K_2^\perp}{K_2}\right)^\vee(h_2)\right) = \{(\delta, p) \in F_B \times B | p \in \Delta\} \subset G_\omega(2, 6) \times G_\omega(3, 6)$$

The projections from I to F_B and B will be denoted by p_1 and p_2 .

2.1 A morphism from F_B to $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$

Each of the 4 conics C_i will enable us to construct a morphism from F_B to $\mathbb{P}_1$. We have the following geometric hint to expect at least a rational map: A general element δ of F_B gives an isotropic 2-dimensional subspace L_δ of W . In general, the projectivisation of $L_\delta^\perp$ meets the plane containing C_i in a point p . There is at least an element m of Δ such that $p \in \pi_m$, so $m \in \Delta \subset B \subset H_i$. Now, the definition of H_i and lemma 1.1 prove that p must be on C_i .

But to show that it is everywhere defined, we will use the vector bundle E_i . We start by constructing line bundles on F_B .

Remark 2.2 Any line Δ included in the hyperplane section $\bar{H}_{u_i} = q_2(Z_i)$ of B intersect the veronese surface V_i . Furthermore, any such line is in a quadric Z_{i,p_i} for a unique point p_i of C_i . So the set $v_i = \{\delta \in F_B | \Delta \subset \bar{H}_{u_i}\}$ is a divisor in F_B .

Proof: As the line Δ is in $\bar{H}_{u_i}$, we have from lemma 1.1 that for any $b \in \Delta$, the plane π_b intersect the conic C_i . The line $\mathbb{P}(L_\delta)$ must intersect C_i in some point p_i . Indeed, if it was not the case, this line would be orthogonal to C_i , so it would be in the plane π_{u_i} , but any line in this plane intersect C_i .

So the line Δ is in the quadric Z_{i,p_i} . Note that $\mathbb{P}(L_\delta) \cap C_i$ can't contain another point, because it would imply that $\Delta \subset V_i$. Furthermore, the proposition 1.9 implies that $Z_{i,p_i} \cap V_i$ is a plane section of the quadric Z_{i,p_i} , so Δ intersect V_i in a single point. $\square$

Remark 2.3 For any point p_i of C_i , the scheme $p_2^{-1}(Z_{i,p_i})$ has several irreducible components of dimension 2, but some of these component are contracted by p_1 to a curve. Denote by A_{i,p_i} the 2 dimensional part of $p_1(p_2^{-1}(Z_{i,p_i}))$.

Proof: The components of $p_2^{-1}(Z_{i,p_i})$ corresponding to the lines included in Z_{i,p_i} are contracted to curves. $\square$

Proposition 2.4 The sheaf $p_{1*}p_2^*E_i$ is a line bundle on F_B . There is a natural map f_i from $H^0(\mathcal{O}_{C_i}(\sigma_i))^\vee \otimes \mathcal{O}_{F_B}$ to the dual bundle of $(p_{1*}p_2^*E_i)$. The image of f_i is also a line bundle on F_B , we will denote it by $\mathcal{O}_{F_B}(\alpha_i)$. By construction, for any $p_i \in C_i$, the divisor A_{i,p_i} will be in the linear system $|\mathcal{O}_{F_B}(\alpha_i)|$.

Proof: By the corollary 1.7, the bundle E_i is a quotient of $6\mathcal{O}_B(-1)$ and by definition 1.2, it is a subsheaf of $2\mathcal{O}_B$. So its restriction to any line Δ included in B must be $\mathcal{O}_\Delta \oplus \mathcal{O}_\Delta(-1)$. So $R^1p_{1*}p_2^*E_i = 0$ and $p_{1*}p_2^*E_i$ is a line bundle. Denote this line bundle by $\mathcal{O}_{F_B}(-\alpha'_i)$. Dualising and twisting the exact sequence defining E_i , we have the following exact sequence:

$$0 \rightarrow L \otimes \mathcal{O}_B(-2h_3) \rightarrow E_i(-h_3) \rightarrow \mathcal{L}_i \rightarrow 0 \quad (2)$$

where $\mathcal{L}_i$ is supported on the hyperplane section $\bar{H}_{u_i}$, and is singular along the veronese surface V_i . As the incidence I is $\text{Proj}((\frac{K_2^\perp}{K_2})^\vee(h_2))$ (where K_2 is the tautological sub-bundle of $W \otimes \mathcal{O}_{G_\omega(2,W)}$), the relative dualising sheaf ω_{p_1} is $\mathcal{O}_I(2h_2 - 2h_3)$. So we have $R^1p_{1*}(p_2^*E_i(-h_3)) = \mathcal{O}_{F_B}(\alpha'_i - 2h_2)$. So the base locus of this pencil of sections of $\mathcal{O}_{F_B}(\alpha'_i)$ is the support of $R^1p_{1*}p_2^*(\mathcal{L}_i)$. We will now prove that this sheaf is a line bundle on v_i .

The morphism p_1 is projective of relative dimension 1. So, for any point δ of F_B and any coherent sheaf F on B , the fiber $(R^1p_{1*}p_2^*F)_\delta$ is $H^1(F \otimes \mathcal{O}_\Delta)$, where Δ is the line in B corresponding to δ . The restriction of the sequence 2 to Δ gives the surjection:

$$2\mathcal{O}_\Delta(-2) \rightarrow \mathcal{O}_\Delta(-1) \oplus \mathcal{O}_\Delta(-2) \rightarrow \mathcal{L}_i \otimes \mathcal{O}_\Delta \rightarrow 0 \quad (3)$$

When the line Δ is not in the hyperplane section $\bar{H}_{u_i}$, the sheaf $\mathcal{L}_i \otimes \mathcal{O}_\Delta$ is supported by the point $\bar{H}_{u_i} \cap \Delta$, so in this case we have $h^1(\mathcal{L}_i \otimes \mathcal{O}_\Delta) = 0$. Now, when the line Δ is in $\bar{H}_{u_i}$, the sheaf $\mathcal{L}_i \otimes \mathcal{O}_\Delta$ has generic rank 1 because the veronese surface V_i can't contain

the line Δ . We have proved in lemma 2.2 that Δ intersect V_i , hence for any element of L , the corresponding section of $E_i(h_3)$ vanishes on Δ , so the map $2\mathcal{O}_\Delta(-2) \rightarrow \mathcal{O}_\Delta(-2)$ induced by the sequence (3) is zero, and for any δ in v_i , we have $h^1(\mathcal{L}_i \otimes \mathcal{O}_\Delta) = 1$, and $R^1 p_{1*} p_2^*(\mathcal{L}_i)$ is a line bundle on v_i .

So we have proved that the base locus of $|(p_{1*}(p_2^*E_i))^\vee|$ is the divisor v_i . In other words, the image of f_i is the line bundle $\mathcal{O}_{F_B}(\alpha_i) = (p_{1*}(p_2^*E_i))^\vee(-v_i)$, which is by construction a quotient of $L \otimes \mathcal{O}_{F_B}$. By definition A_{i,p_i} is the closure of $\{\delta \in F_B \mid \text{lenght}(\Delta \cap Z_{i,p_i}) = 1\}$ which was identified set theoretically with an element of the linear system $|\alpha_i|$, so we conclude the proof with lemma : $\square$

Lemma 2.5 *For a generic choice of a point p_i on C_i , the support of the sheaf $R^1 p_{1*}(p_2^* I_{Z_{i,p_i} \cup V_i}(-h_3))$ represent the class α'_i , and all its irreducible components are reduced.*

Proof: First notice that the point p_i on C_i , gives a section of $E_i(h_3)$, so an exact sequence:

$$0 \rightarrow \mathcal{O}_B(-2h_2) \rightarrow E_i(-h_3) \rightarrow I_{Z_{i,p_i} \cup V_i}(-h_3) \rightarrow 0$$

which gives a section of $\mathcal{O}_{F_B}(\alpha'_i)$ vanishing on the support of the sheaf $R^1 p_{1*}(p_2^* I_{Z_{i,p_i} \cup V_i}(-h_3))$. But this is the definition in [G-P] of the scheme structure on the set of lines included in B and intersecting $Z_{i,p_i} \cup V_i$. So to show that this scheme structure is reduced on each component, we have to prove that Z_{i,p_i} and V_i are not in the ramification of the morphism: $p_2 : I \rightarrow B$. But a general point m on $Z_{i,p_i} \cap V_i$ is the intersection of Z_{i,p_i} and another quadric Z_{i,p'_i} , so there pass 4 distinct lines through m , and from remark 2.2 there are no other lines in B through m . So m is not in the ramification of the morphism $p_2 : I \rightarrow B$ because it is of degree 4. (Cf lemma 3.1). $\square$

2.2 Description of the morphism

In the previous section, we have constructed 4 morphism f_i from F_B to $\mathbb{P}_1$. We will now prove that the morphism from F_B to $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ is an embedding for a generic B , and that its image is an hyperplane section.

NOTATIONS:

For a point p_i in the conic C_i , we denote by $\mathcal{C}_B$ the affine cone over B . We consider:

$$F_{p_i} = \{\delta \in G_\omega(2, W) \mid \delta \wedge p_i \in \mathcal{C}_B\}$$

Unfortunately, we have to remark that $f_i^{-1}(p_i)$ is not exactly $F_{p_i} \cap F_B$:

Remark 2.6 *The intersection $F_{p_i} \cap F_B$ is equal to $G_\omega(2, p_i^\perp) \cap F_B$. It contains $f_i^{-1}(p_i)$ and residual curves corresponding to the lines included in the quadric Z_{i,p_i} .*

Proof: If δ is already an element of F_B , then any isotropic plane of $\mathbb{P}(W)$ containing the line $\mathbb{P}(L_\delta)$ is an element of B (Cf remark 2.1). For $\delta \in G_\omega(2, p_i^\perp)$, the plane $\pi_{\delta \wedge p_i}$ is isotropic, so we have $\delta \wedge p_i \in \mathcal{C}_B$ and $G_\omega(2, p_i^\perp) \cap F_B = F_{p_i} \cap F_B$. Now, remark that in

$G_\omega(2, W)$ the scheme $G_\omega(2, p_i^\perp)$ is the zero locus of a section of $K_2^\vee$. The restriction of this section to F_B vanishes on the divisor A_{i,p_i} defined in proposition 2.4, so $F_B \cap F_{p_i}$ contains a divisor of class α_i and the zero locus of a section of $(K_2^\vee)_{|F_B}(-\alpha_i)$ corresponding to the lines includes in Z_{i,p_i} . $\square$

Nevertheless, we have the following:

Lemma 2.7 *For the generic double hyperplane section B of $G_\omega(3, W)$, we can find points p_i (resp p_j) in the conics C_i (resp C_j) such that $F_{p_i} \cap F_{p_j}$ is a smooth conic in $G_\omega(2, W)$. Furthermore, this conic is in F_B and represent the class $\alpha_i \cdot \alpha_j$.*

Proof: We can choose p_i and p_j respectively in C_i and C_j such that, p_i is not in $p_j^\perp$. Remark first that this implies that the intersection $F_{p_i} \cap F_{p_j}$ is automatically included in F_B because p_i and p_j are never in the same isotropic plane. This also implies that the isotropic grassmannian $G_\omega(2, p_i^\perp \cap p_j^\perp)$ is a smooth 3-dimensional quadric. The intersection $F_{p_i} \cap F_{p_j}$ is given in this grassmannian by the 2 conditions: $d \wedge p_i \in H_j$ and $d \wedge p_j \in H_i$ for an element d of $G_\omega(2, p_i^\perp \cap p_j^\perp)$. Indeed, as $d \wedge p_i$ and $d \wedge p_j$ represent isotropic planes, the conditions $d \wedge p_i \in H_i$ and $d \wedge p_j \in H_j$ are automatically satisfied because $p_i \in C_i$ and $p_j \in C_j$ (Cf lemma 1.1). Let δ be the intersection of $p_j^\perp$ and the 3 dimensional vector space U_i represented by the contact point u_i of H_i . The space δ is an element of $G_\omega(2, p_i^\perp \cap p_j^\perp)$ such that $p_i \wedge \delta$ is not in H_j (because it is u_i , and B is smooth) but $p_j \wedge \delta \in H_i$ according to lemma 1.1. So the two hyperplane sections are independent and we have proved that $F_{p_i} \cap F_{p_j}$ is a (may be singular) conic.

Note that from the genericity of B we could assume also that $\mathbb{P}(p_i^\perp) \cap C_j = \{a_1, a_2\}$ and $\mathbb{P}(p_j^\perp) \cap C_i = \{b_1, b_2\}$ are 4 distinct points. According to lemma 1.1, the lines (p_i, a_1) , (p_i, a_2) , (p_j, b_1) , (p_j, b_2) are in $F_{p_i} \cap F_{p_j}$, but no three of those lines are in the same plane, so the conic $F_{p_i} \cap F_{p_j}$ contains 4 points with no trisecant line. Hence the conic must be smooth. To conclude that this conic represent the class $\alpha_i \cdot \alpha_j$, we just have to prove that the residual curve of $F_{p_i} \cap F_B$ don't intersect F_{p_j} . But if δ is such that $\Delta \subset Z_{i,p_i}$, the line $\mathbb{P}(L_\delta)$ contains the point p_i which is not orthogonal to p_j , so $\delta \notin F_{p_j}$. $\square$

Lemma 2.8 *When i, j, k are distinct, the morphism from F_B to $C_i \times C_j \times C_k$ is dominant.*

Proof: According to lemma 2.7, for a generic choice of $p_i \in C_i$ and $p_j \in C_j$ the subvariety $F_{p_i} \cap F_{p_j}$ of F_B is a smooth conic, and we can also assume that the line $(p_i p_j)$ doesn't intersect C_k . Assume that the induced map from $F_{p_i} \cap F_{p_j}$ to C_k is not dominant, then there is a point $p_k \in C_k$ such that $F_{p_i} \cap F_{p_j} \subset F_{p_k}$. So for any element d of $F_{p_i} \cap F_{p_j}$ the corresponding line D is in the plane $\mathbb{P}(\langle p_i, p_j, p_k \rangle^\perp)$. As $p_k \notin (p_i p_j)$, the vector space $\langle p_i, p_j, p_k \rangle^\perp$ has dimension 3, and this contradicts the fact that $F_{p_i} \cap F_{p_j}$ is a smooth conic. $\square$

At this point, we need more details about the embedding of F_B in $G_\omega(2, W)$:

NOTATIONS:

Still denote by K_2 the tautological rank 2 subsheaf of $W \otimes \mathcal{O}_{G_\omega(2, W)}$, and by $-h_2$ and c_2 its first and second Chern classes. As $G_\omega(2, W)$ is an hyperplane section of $G(2, W)$, we will do the computations in $G(2, W)$.

Remark 2.9 *The Chow ring of $G(2, 6)$ is*

$$\mathbb{C}[h_2, c_2]/(h_2^5 + 3h_2c_2^2 - 4h_2^3c_2, -h_2^4c_2 + 3h_2^2c_2^2 - c_2^3)$$

We have:

Lemma 2.10 *The variety F_B is obtained in $G_\omega(2, W)$ as the zero locus of a section of the bundle $(\frac{K_2^\perp}{K_2})^\vee(h_2) \oplus (\frac{K_2^\perp}{K_2})^\vee(h_2)$. For a general choice of B , F_B is smooth with $\omega_{F_B} = \mathcal{O}_{F_B}(-h_2)$. The class of F_B in the chow ring of $G_\omega(2, W)$ is $4 \cdot (h_2^2 - c_2)^2$, and F_B has degree 24 in $G_\omega(2, W)$. In the Chow ring of F_B , we have the extra relation: $h_2^2 = 3c_2$.*

Proof: To compute Chern classes, we just have to remark that $K_2^\perp$ is isomorphic to the dual of the tautological quotient. The vanishing locus claim is a consequence of the fact that B is a double hyperplane section of $G_\omega(3, W)$, and that we have from the definition of I the equality: $p_{1*}(p_2^*\mathcal{O}_{G_\omega(3, W)}(h_3)) = (\frac{K_2^\perp}{K_2})^\vee(h_2)$. (with the notations introduced in §2). So the choice of a generic 2-dimensional subspace of $H^0(\mathcal{O}_{G_\omega(3, W)}(h_3))$ correspond to the choice of a generic section of $(\frac{K_2^\perp}{K_2})^\vee(h_2) \oplus (\frac{K_2^\perp}{K_2})^\vee(h_2)$. Hence, for a general choice of B , F_B will be smooth because $K_2^\perp(h_2)$ is globally generated, and so is $\frac{K_2^\perp}{K_2}(h_2) \simeq (\frac{K_2^\perp}{K_2})^\vee(h_2)$. We can now conclude with the computations using the remark 2.9. $\square$

Lemma 2.11 *In the Chow ring of F_B we have $\alpha_i^2 = 0$, $\alpha_i \cdot \alpha_j \cdot h_2 = 2$ and $\alpha_i \cdot \alpha_j \cdot \alpha_k = 1$ when i, j, k are distinct. (where h_2 is the hyperplane section of F_B)*

Proof: In proposition 2.4 we already proved that $\alpha_i^2 = 0$. We have proved in the lemma 2.8, that for a generic choice of p_i, p_j, p_k the intersection $F_{p_i} \cap F_{p_j} \cap F_{p_k}$ is not empty. Furthermore, this intersection is included in the smooth conic $F_{p_i} \cap F_{p_j}$ and the line $G_\omega(2, p_i^\perp \cap p_j^\perp \cap p_k^\perp)$. So it must be a point because the intersection of a smooth conic and a line in the 3-dimensional smooth quadric $G_\omega(2, p_i^\perp \cap p_j^\perp)$ must be empty or a point. So $\alpha_i \cdot \alpha_j \cdot \alpha_k = 1$. $\square$

Lemma 2.12 *The hyperplane section h_2 of F_B is linearly equivalent to $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$.*

Proof: We deduce from the lemma 2.11 that the image of the morphism from F_B to $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ is a divisor $\bar{F}$ of class $(1, 1, 1, 1)$. Furthermore, the lemmas 2.8 and 2.11 imply that $\psi : F_B \rightarrow \bar{F}$ is a birational morphism.

Now we should notice that $\bar{F}$ is normal because the restriction of $\psi : F_B \rightarrow \bar{F}$ to a double hyperplane section of $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ is not ramified. Indeed, from the lemma 2.11 we have: $(\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4)^3 = 24 = h_2 \cdot (\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4)^2$, so the restriction of ψ to a general double hyperplane section of $\bar{F}$ is not ramified.

From the normality of $\bar{F}$, we obtain a map $\psi^*(\omega_{\bar{F}}) \rightarrow \omega_{F_B}$. This implies that the divisor $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 - h_2$ of F_B is effective in F_B . To obtain the result, we will show that its intersection with h_2^2 is zero. Remarking from lemma 2.10 that on F_B we have $h_2^3 = 24$, we will end the proof after the following lemma:

Lemma 2.13 *The image of the class α_i in the Chow ring of $G_\omega(2, W)$ is $h_2.c_2.2(h_2^2 - c_2)$, and in the Chow ring of F_B we have $\alpha_i.h_2^2 = 6$.*

Proof: First, chose an hyperplane H' such that $B = G_\omega(3, W) \cap H' \cap H_i$. Let $F_{H'}$ and F_{H_i} be the variety of lines included respectively in $\bar{H}'$ and $\bar{H}_i$, and denote by Y the Grassmannian $G_\omega(2, p_i^\perp)$. We proved in remark 2.6 that α_i is represented by the 2-dimensional part of $Y \cap F_B$ for some point p_i on the conic C_i . From the definition of B , we have $F_B = F_{H'} \cap F_{H_i}$. The variety $F_{H'}$ and F_{H_i} represent the class $c_2((\frac{K_2^\perp}{K_2})^\vee(h_2))$ in the Chow ring of $G_\omega(2, W)$, but the intersection $Y \cap F_{H_i}$ has codimension one in Y . Indeed, for any l in Y the point $p_i \wedge l$ is already in $\bar{H}_i$. The restriction of the sheaf $(K_2)^\perp$ to Y has a section given by p_i , and the intersection $G_\omega(2, p_i^\perp) \cap F_{H_i}$ is the vanishing locus of a section of $\frac{K_{2|Y}^\perp}{p_i + K_{2|Y}}(h_2)$. Remarking that Y represent the class c_2 in $G_\omega(2, W)$, we conclude that the class α_i is equivalent to $h_2.c_2.c_2((\frac{K_2^\perp}{K_2})^\vee(h_2))$ in the Chow ring of $G_\omega(2, W)$.

Using the remark 2.9 we compute that $h_2^2.\alpha_i = 6$. This ends the proof of lemma 2.13, and also of lemma 2.12. $\square$

In conclusion, we have the following:

Theorem 2.14 *The variety of lines included in a generic double hyperplane section of $G_\omega(3, 6)$ is an hyperplane section of $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$.*

Proof: In lemma 2.12 we proved that $h_2 \sim \sum_{i=1}^4 \alpha_i$, but we have $h^0(\mathcal{O}_{\bar{F}}(1, 1, 1, 1)) = 15$ and $h^0(\mathcal{O}_{G_\omega(2, W)}(h_2)) = 14$, so the morphism $\psi : F_B \rightarrow \bar{F}$ is an isomorphism. $\square$

3 The Chow ring of B

We keep the notations of section 2. We will study here the Chow ring of B . The 4-dimensional variety B is a generic double hyperplane section of $G_\omega(3, 6)$. Its Chow ring in codimension 2 will appear to be surprisingly bigger than the codimension 2 part of the Chow ring of $G_\omega(3, 6)$. To study it, we will first compare it to the Chow ring of I via the projection $p_2 : I \rightarrow B$.

Lemma 3.1 *The variety B contains 16 quadric cones of dimension 2. For any point m different from the 16 vertex of those cones, the fiber $p_2^{-1}(\{m\})$ has length 4.*

Proof: Let m be any point of B . The lines containing m and included in B are the lines containing m and included in the tangent cone of B . The tangent cone to $G_\omega(3, W)$ is a cone over a veronese surface. The smoothness of B at m implies that the tangent cone of B at m is a cone over the intersection of a veronese surface by a $\mathbb{P}_3$. So it is either 4 lines or a 2 dimensional quadric cone with smooth basis.

Now, let Γ be such a cone included in B , we can deduce from lemma 3.5 of [I] that there is a point e of $\mathbb{P}(W)$ included in all the planes π_u for u in Γ . So Γ is the intersection of the 3-dimensional quadric $Q_e = \{e \wedge l \mid l \in G_\omega(2, \frac{e^\perp}{e})\}$ with 2 hyperplanes containing B . This proves that one of them contains Q_e . We deduce from the proposition 3.3 of

[I] that this hyperplane must be one of the H_i defined in section 1 and that e is on one of the four conics C_i . But the description of the incidences Z_i in proposition 1.9 proves that there are only 4 cones for each C_i . $\square$

NOTATIONS:

*In the sequel, we will denote the Chow ring of a variety X by A_X . Let $c'_1 = h_3$, c'_2 and c'_3 be the Chern classes of the tautological quotient Q_3 of $W \otimes \mathcal{O}_{G_\omega(3,W)}$. Denote by h'_3 the class $p_2^*h_3$ in A_I , and let a_i be the second Chern class of the bundle E_i .*

First recall from the section 2 and the lemma 2.10, that the Chow ring of the incidence I is:

Lemma 3.2 *The Chow ring of I is*

$$A_I = \frac{A_{F_B}[h'_3]}{(h_3'^2 - 2.h_2.h'_3 + \frac{4}{3}.h_2^2)}$$

So we have to get more informations on F_B .

Lemma 3.3 *For each choice of i , the variety F_B can be identified with the blow up of $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ in an elliptic sextic curve, where v_i is the exceptionnal divisor.*

The choice of one of the $\mathbb{P}_1$ (ie: the marking of one of the conics C_i) in theorem 2.14 enable us to consider the variety F_B as an incidence:

$$\{(x, h) | x \in \mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1, h \in L, x \in h\}$$

where L is a marked subspace of $H^0(\mathcal{O}_{\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1}(1, 1, 1))$. In other words, for each i , the variety F_B with le line bundle $h_2 - \alpha_i$ is identified with the blow up of $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ in an elliptic sextic curve. By construction $|\alpha_i|$ is the marked pencil of hyperplanes.

To show that v_i is the exceptionnal divisor, we need to prove that $2\alpha_i + v_i = h_2$. Take any subset $\{i, j, k\}$ of $\{1, 2, 3, 4\}$ of cardinal 3, and chose a point p_j on C_j and another one p_k on C_k such that $\mathbb{P}(p_j^\perp \cap p_k^\perp) \cap C_i = \emptyset$. The divisor v_i was constructed in proposition 2.4 as the set of lines included in the hyperplane section $\bar{H}_{u_i}$ of B . So $\delta \in v_i$ implies that $\Delta \subset \bar{H}_{u_i}$ which implies that the line $\mathbb{P}(L_\delta)$ intersect C_i , so it can't be in $\mathbb{P}(p_j^\perp \cap p_k^\perp)$, and we have $\alpha_j \cdot \alpha_k \cdot v_i = 0$. So set theoretically, v_i is the exceptionnal divisor, but we proved in lemma 2.5 that v_i is reduced. $\square$

Lemma 3.4 *The projection $p_{1*}(p_2^*a_i)$ of the second Chern class of $p_2^*E_i$ to A_{F_B} is $h_2 - \alpha_i$. Furthermore, we have in A_I the equality: $p_2^*a_i = h'_3 \cdot (h_2 - \alpha_i) + h_2(2\alpha_i - h_2)$*

Proof: We know from the lemma 3.3 and proposition 2.4, that the first Chern class of $p_{1*}(p_2^*E_i)$ is $-\alpha_i - v_i = \alpha_i - h_2$. So the first assertion is a direct consequence of Riemman-Roch-Grothendieck's theorem because $(p_2^*E_i)$ have no higher direct images by p_1 . Indeed, we can compute in A_I using the equality: $\omega_{p_1} = \mathcal{O}_I(2h_2 - 2h_3)$ obtained in §2, and it gives that the first Chern class of $p_{1*}(p_2^*E_i)$ is $-p_{1*}p_2^*a_i$.

The second assertion is obtained by the evaluation map: $p_1^*p_{1*}p_2^*E_i \rightarrow p_2^*E_i$. We can compute its cokernel as we done in lemma 1.10, but the vanishing of $R^1p_{1*}(p_2^*E_i)$ gives the following exact sequence:

$$0 \rightarrow p_1^*p_{1*}p_2^*E_i \rightarrow p_2^*E_i \rightarrow \mathcal{O}_I(-h'_3) \otimes p_1^*R^1p_{1*}(p_2^*(E_i(-h_3))) \rightarrow 0$$

so we have by relative duality the extention:

$$0 \rightarrow \mathcal{O}_I(p_1^*\alpha_i - p_1^*h_2) \rightarrow p_2^*E_i \rightarrow \mathcal{O}_I(p_1^*h_2 - p_1^*\alpha_i - h'_3) \rightarrow 0 \quad (4)$$

which gives the computation of $p_2^*a_i$. $\square$

Lemma 3.5 *Let $\gamma \in A_B^2$, then the class $p_2^*\gamma$ can be written in A_I by $h'_3.\gamma_0 + \gamma_1$ with γ_i in $A_{F_B}^{i+1}$, and where γ_i is in the vector space generated by $h_2^i\alpha_1, \dots, h_2^i\alpha_4$. More precisely, we have: $2h_2.\gamma_0 + \gamma_1 \in \mathbb{Q}.h_2^2$*

Proof: We can first find classes γ_i in $A_{F_B}^{i+1}$ such that $p_2^*\gamma = h'_3.\gamma_0 + \gamma_1$. Now remark that A_B^3 is one dimensional, so $h_3.\gamma$ is propotional to h_3^3 . But from the relation $h_3'^2 = 2.h_2.h_3' - \frac{4}{3}.h_2^2$, the class $p_{1*}h_3'^3$ is proportional to h_2^2 . So the class $p_{1*}p_2^*\gamma$ is also in $\mathbb{Q}.h_2^2$, and we have $2h_2.\gamma_0 + \gamma_1 \in \mathbb{Q}.h_2^2$. We can now conclude with the description of F_B in lemma 3.3 that γ_1 and h_2 are in the vector space generated by $\alpha_1, \dots, \alpha_4$, which gives the lemma. $\square$

Lemma 3.6 *The classes (a_1, a_2, a_3, a_4) form a basis of the vector space A_B^2 . We have in A_B the relation $2(a_1 + a_2 + a_3 + a_4) = 3h_3^2$.*

Proof: From the lemma 3.1, the map $p_2^* : A_B^2 \rightarrow A_I^2$ is injective, so from the lemma 3.5, A_B^2 is generated by the set of classes $\{p_{2*}(h'_3.\alpha_i), p_{2*}(h_2\alpha_i)\}_{i=1\dots 4}$. As the picard group of B is generated by h_3 , all the classes $p_{2*}(h'_3.\gamma_0)$ are proportional to h_3^2 . Now, we use the relation $\frac{4}{3}.h_2^2 = 2.h_2.h_3' - h_3'^2$ to eliminate h_2^2 in the expression of $p_2^*a_i$ found in lemma 3.4. So, we obtain that a_i is in the affine space $\frac{1}{2}.p_{2*}(h_2\alpha_i) + \mathbb{Q}.h_3^2$. So we have proved that A_B^2 is generated by $h_3^2, a_1, a_2, a_3, a_4$. Furthermore, as we have $p_{1*}p_2^*a_i = h_2 - \alpha_i$, which is a free family in $A_{F_B}^1$, the family $(a_1, \dots, a_4)$ is free in A_B^2 .

To obtain the relation with h_3^2 , we substitute the expression of $p_2^*a_i$ of lemma 3.4 in the relation found in lemma 2.12. We eliminate $h'_3.h_2$ with the relation $\frac{4}{3}.h_2^2 + h_3'^2 = 2.h_2.h_3'$, and we obtain: $\sum_{i=1}^4 p_2^*a_i = \frac{3}{2}h_3'^2$. $\square$

So we are now ready to describe the Chow ring of B .

Proposition 3.7 *The Chow ring of B is $\mathbb{Q}[h_3, a_1, a_2, a_3, a_4]/\mathcal{I}$ where $\mathcal{I}$ is generated by $3h_3^2 - 2.\sum_{i=1}^4 a_i$, $(8h_3.a_i - 3h_3^3)_{i \in \{1, \dots, 4\}}$, $(8.a_i.a_j - h_3^4)_{i \neq j, (i,j) \in \{1, \dots, 4\}^2}$. (The class of a point is $\frac{a_1.a_2}{2}$, the class of the veronese V_i is $2a_i - \frac{1}{2}h_3^2$, and $[V_i]^2$ is 4 points).*

Proof: As A_B^i is known to be one dimensional for $i \in \{0, 1, 3, 4\}$, we just need to compute the relations, and it can be done by calculating the degrees, because we have found the structure of A_B^2 in lemma 3.6. The relations $(8h_3.a_i - 3h_3^3)$ are consequences of the degree of $G_\omega(3, 6)$ (ie 16) and the fact that a_i can be represented by the union of a quadric Z_{i,p_i} and the veronese surface V_i .

Now remark that the class of V_i is in the vector space generated by a_i and h_3^2 . Indeed, we have $p_2^*[V_i] = h_3'.v_i + \gamma_1$ and the lemmas 3.3 and 3.5 give $v_i = h_2 - 2\alpha_i$ and $\gamma_1 \in h_2^2.\mathbb{Q} + h_2\alpha_i.\mathbb{Q}$, so $[V_i]$ is in the vector space generated by h_3^2 and a_i , and then $[Z_{i,p_i}]$ also. Computing their degree, we have: $[V_i] = 2a_i - \frac{1}{2}h_3^2$ and $[Z_{i,p_i}] = \frac{1}{2}h_3^2 - a_i$. The last relations are then consequences of the fact that we have: $[V_i].[V_j] = 0$ for $i \neq j$. $\square$

NB:

It could be usefull to have the link with the ring of $G_\omega(3, 6)$ and the Chern classes of Q_3 , so we state the following:

Remark 3.8 *Still denote by c'_i the Chern classes of the tautological quotient Q_3 . The Chow ring of $G_\omega(3, 6)$ is*

$$\mathbb{Q}[c'_1, c'_2, c'_3]/(c_3'^2, c_2'^2 - 2c_1'c_3', c_1'^2 - 2c_2')$$

On B we have the additional relation $c'_1c'_2 = 4c'_3$. In particular the rank of $A_{G_\omega(3,6)}^2$ is only one.

Proof: The Chow ring of $G_\omega(3, 6)$ is given by

$$\mathbb{Q}[x, y, z]/((xyz)^2, x^2y^2 + y^2z^2 + z^2x^2, x^2 + y^2 + z^2)$$

where $c'_1 = x + y + z$, $c'_2 = xy + yz + zx$ and $c'_3 = xyz$. $\square$

4 Application to quadratic normality

In this part, we explain the link with the congruence of lines found in [M-dP]. They started with the intersection Γ of $G(2, 6)$ by a very particular $\mathbb{P}_{11}$ (NB: it was proved in [M-M], that the choice of this $\mathbb{P}_{11}$ is unique up to the action of GL_6). Then they chosed a general quadric containing a fixed subscheme of Γ (which contains the singular locus of Γ) to obtain a reducible intersection with Γ . Then they checked with Macaulay2 that one of these irreducible components is smooth of degree 16 and sectional genus 9.

Here we prove that the generic Fano 4-fold of genus 9 can be obtained by their construction, and that the choice we need to do is generically finite. More precisely, the choice of Γ correspond to the choice of a tangent hyperplane H to $G_\omega(3, 6)$, and the choice of the quadric will correspond to the choice of a non zero element in $|\mathcal{O}_{\bar{H}}(h_3)|$.

Proposition 4.1 *For any choice of i , the bundle $E_i(h_3)$ gives an embedding of the Fano manifold B in the grassmannian $G(2, 6)$ as the congruence of lines constructed in [M-dP] (Theorems 8,9,10).*

Proof: Their description of the congruence is in terms of equations, so we need to make an adapted choice of coordinates to get the link. Choose a basis $w_0, \dots, w_5$ of W such that $\omega^\vee = w_0 \wedge w_3 + w_1 \wedge w_4 + w_2 \wedge w_5$. Let A and B be the vector spaces generated respectively by w_0, w_1, w_2 and w_3, w_4, w_5 , in particular, the form ω gives an identification between $A^\vee$ and B . The decomposition $W = A \oplus B$ gives a decomposition of $\bigwedge^3 W$. So

we can represent an element of $\bigwedge^3 W$ as in [I] by (a, X, Y, b) , with $a \in \bigwedge^3 A$, $b \in \bigwedge^3 B$, $X \in \text{Hom}(A, B)$, $Y \in \text{Hom}(B, A)$. The equations of $G(3, 6)$ are as: $\bigwedge^2 X = aY$, $\bigwedge^2 Y = bX$, $YX = ab.I_3$, and to obtain $G_\omega(3, 6)$ we had the linear relations $X = {}^tX$ and $Y = {}^tY$, so we take $X = \begin{pmatrix} x_0 & x_1 & x_2 \\ x_1 & x_5 & x_3 \\ x_2 & x_3 & x_4 \end{pmatrix}$ and $Y = \begin{pmatrix} y_0 & y_1 & y_2 \\ y_1 & y_5 & y_3 \\ y_2 & y_3 & y_4 \end{pmatrix}$. (Cf [I]).

Now we need to choose an hyperplane H as in section 1.1. So we consider the hyperplane $y_5 = y_2$. It is tangent to $G_\omega(3, 6)$ in $w_0 \wedge w_1 \wedge w_2$ and contains $w_3 \wedge w_4 \wedge w_5$. Furthermore, the conic described in lemma 1.1 is parametrized by $\lambda^2 w_0 + \lambda \mu w_1 + \mu^2 w_2$, so the incidence Z_H is given in $\mathbb{P}_1 \times \bar{H}$ by the equations:

$$a = 0, X \cdot \begin{pmatrix} \lambda^2 \\ \lambda \mu \\ \mu^2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -\mu & \lambda & 0 \\ 0 & -\mu & \lambda \end{pmatrix} \cdot Y = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

The 6 dimensional vector space $H^0(\mathcal{I}_{Z_H}(1, 1))$ is generated by

$$(-a.\lambda, -a.\mu, \lambda y_1 - \mu y_0, \lambda y_2 - \mu y_1, \lambda y_3 - \mu y_2, \lambda y_4 - \mu y_3)$$

So, the sheaf $\mathcal{E}$ constructed in proposition 1.2 is the image of the map:

$$6\mathcal{O}_{\bar{H}}(-1) \xrightarrow{\begin{pmatrix} a & 0 & y_0 & y_1 & y_2 & y_3 \\ 0 & a & -y_1 & -y_2 & -y_3 & -y_4 \end{pmatrix}} 2\mathcal{O}_{\bar{H}}$$

which is exactly the map of [M-dP] th 9. Now, a generic hyperplane section of $G_\omega(3, 6)$ gives a linear relation between $a, (x_i), (y_i), b$ which is exactly the relation 29 of [M-dP], which gives the identification with their congruence of lines. $\square$

So we can remark that the computations they made in affine charts of $G(2, 6)$ can be globally done in $G_\omega(3, 6)$.

4.1 Geometry of the focal locus of Mezzeti-de Poi's congruence

Here we chose one of the 4 bundles, say E_1 . Denote by h the bundle $\mathcal{O}_{\text{Proj}(E_1(h_3))}(1)$. We proved in lemma 1.7 that the linear system $|h|$ gives a morphism from $\text{Proj}(E_1(h_3))$ to $\mathbb{P}_5$.

NOTATIONS:

Denote by r the above projection from $\text{Proj}(E_1(h_3))$ to $\mathbb{P}_5$, and by p_B the projection from $\text{Proj}(E_1(h_3))$ to its basis B .

Following [M-dP], the focal locus of Mezzeti-de Poi's congruence is defined like this:

Lemma 4.2 *The morphism from $r : \text{Proj}(E_1(h_3)) \rightarrow \mathbb{P}_5$ is birational. The class of the exceptional divisor $\mathcal{R}$ in $\text{Proj}(E_1(h_3))$ is $4h - h_3$.*

Proof: The degree of r is given by the length of the degeneracy locus of a generic map $5\mathcal{O}_B \rightarrow E_1(h_3)$, so it is the fourth segre class of E_1 , which is the class of one point from proposition 3.7. So r is birational, and the class of the exceptional divisor follows. $\square$

The manifold B gives the family $LB = \{\mathbb{P}(E_{1,u}) | u \in B\}$ of lines in $\mathbb{P}_5$. From the lemma 4.2, those lines are quadrisecant to X .

NOTATIONS:

Denote by X the focal locus of B , (ie $X = r(\mathcal{R})$). Recall from [M-dP] that X has dimension 3, degree 6, and is singular along a rational smooth cubic curve C .

From the lemma 4.2, any line of this family intersects X in length 4 or is included in X . We can now describe easily the normalisation $\tilde{X}$ of X , and note that we have as in the Palatini case some kind of duality between X and a family of plane cubics included in X , but here it breaks over the singular locus of X :

Proposition 4.3 *The focal locus X of Mezzeti-de Poi's congruence is a projection of $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ from a line. The variety of pencils of lines belonging to LB is $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ blown up in the elliptic curve of degree 6 which is the double cover of the cubic curve C .*

Proof: We pullback the situation to the incidence point/lines of B via the projection $p_2 : I \rightarrow B$. We still denote by p_2 the projection from $Proj(p_2^*(E_1(h_3)))$ to $Proj(E_1(h_3))$. In the proof of lemma 3.4, we noticed the extension (4), which gives a surjection:

$$p_2^*(E_1(h_3)) \twoheadrightarrow \mathcal{O}_I(p_1^*h_2 - p_1^*\alpha_1)$$

This surjection gives an embedding of I in $Proj(p_2^*(E_1(h_3)))$. The restriction of $p_2^*(h)$ to I is $p_1^*(h_2 - \alpha_1)$, but it is also $p_1^*(\alpha_2 + \alpha_3 + \alpha_4)$ by lemma 2.12. So the linear system $|p_2^*(h)|$ contracts the fibers of $p_1 : I \rightarrow F_B$, and the image of I coincide with the image of F_B by $\alpha_2 + \alpha_3 + \alpha_4$. In conclusion, the map r contracts the divisor $p_2(I)$ to the projection of $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ from a line. $\square$

We can also obtain a description of the plane curves in X related to this family of lines:

Remark 4.4 *Any point of F_B gives a plane in $\mathbb{P}_5$ intersecting X in a point and a plane cubic. Those plane cubics are all singular in a point of C . Only 12 lines included in X are in LB .*

Proof: A point $\delta \in F_B$ correspond to a line Δ included in B . We already proved in §2 that the restriction of E_1 to Δ is always $\mathcal{O}_\Delta \oplus \mathcal{O}_\Delta(-1)$, so the lemma 4.2 implies that the intersection of $\mathcal{R}$ with $p_B^{-1}(\Delta)$ is given by a section of $(S_4(E_1))(3h_3)$. Hence this intersection contains the exceptionnal divisor of $\mathbb{P}(\mathcal{O}_\Delta \oplus \mathcal{O}_\Delta(1))$, so the plane $r(p_B^{-1}(\Delta))$ intersect X in a cubic curve and a unique residual point included in all the lines $r(\mathbb{P}(E_{1,p}))$ for any point p of Δ . Note that the exceptional divisor $\mathcal{R}$ is the image of the composition:

$$I \subset Proj(p_2^*(E_1(h_3))) \rightarrow Proj(E_1(h_3)) \quad (5)$$

so from lemma 3.1 the projection $\mathcal{R} \rightarrow B$ is finite of degree 4 except over 16 points of B , but we will notice later that 4 of them are contracted to points on X . To understand why the plane cubic described above is singular, we first notice that it is the image of the following curve T_Δ in F_B : The closure in F_B of the lines included in B intersecting Δ and different from Δ . A general Δ , intersect the hyperplane section $\bar{H}_{u_1}$ in a single point b which is not on the veronese V_1 . From the corollaries 1.12 and 1.13, there is a single quadric Z_{1,p_1} containing b . The 2 lines containing b and included in this quadric

are included in $\bar{H}_{u_1}$. So, from remark 2.2, they correspond in F_B to points on the exceptional divisor $v_1 \subset F_B$. In conclusion, for a general Δ , the curve T_Δ intersects the exceptional divisor v_1 in 2 points. The image of those 2 points of F_B must be a single point of C because they are in the line $r(\mathbb{P}(p_B^{-1}(b)))$, and this line intersects C in at most one point because b is not in the veronese V_1 . So the plane cubic is singular at this point of C .

For the same reason, if b is one of the 4 vertex of the cones in $\bar{H}_{u_1} \cap B$, $r(\mathbb{P}(p_B^{-1}(b)))$ is contracted to a point on the curve C , that's why only 12 lines of X belong to the family LB . $\square$

As all lines in those planes are trisecant to X , we have from the above remark that:

Remark 4.5 *The triple locus of the projection of $\mathbb{P}_1 \times \mathbb{P}_1 \times \mathbb{P}_1$ from a generic plane is reducible, because it contains the union of 4 lines with a common point.*

We give now some details on the marked “virtual” section related to the anormality of X .

Remark 4.6 *The ramification of the morphism $p_2 : I \rightarrow B$ is $p_1^{-1}(\Sigma)$ for some surface Σ in F_B . The canonical sheaf of Σ is $\omega_\Sigma = \mathcal{O}_\Sigma$, and Σ contains the 16 rational curves parameterizing the 16 cones of B . The image of Σ into X is a section of $\mathcal{O}_X(2)$ that is not in a quadric of $\mathbb{P}_5$.*

Proof: The canonical divisor of I can be computed from §2, and it gives that $\mathcal{R} \sim p_1^*h_2$. The 16 contracted curves must be in $\mathcal{R}$, so we have only to prove that the image of Σ in X is not in a quadric of $\mathbb{P}_5$. But the answer is general for those anormality constuctions: As $\mathcal{R}$ is a divisor in the projective bundle $Proj(E_1(h_3))$, we have the exact sequence:

$$0 \rightarrow \omega_{p_B} \rightarrow \omega_{p_B}(\mathcal{R}) \rightarrow \omega_{\mathcal{R}} \otimes \omega_B^\vee \rightarrow 0$$

where ω_{p_B} is the relative dualising sheaf of $p_B : Proj(E_1(h_3)) \rightarrow B$. The ramification of the restriction of p_B to $\mathcal{R}$ is the zero locus of a section of $\omega_{\mathcal{R}} \otimes \omega_B^\vee$ which gives a non zero element of $H^1(\omega_{p_B}) = \mathbb{C}$. But we saw in the proof of remark 4.4 that the map $I \rightarrow B$ factors through $\mathcal{R}$ (Cf the composition (5)), so we have the result because the above sequence is also:

$$0 \rightarrow \omega_{p_B} \rightarrow \mathcal{O}_{Proj(E_1(h_3))}(2h) \rightarrow \mathcal{O}_{\mathcal{R}}(2h) \rightarrow 0$$

$\square$

4.2 The rank 2 reflexive sheaf on $\mathbb{P}_5$

If we take a general double hyperplane section of X , we obtain a smooth elliptic curve of degree 6. So from Serre's construction, it is the vanishing locus of a section of a rank 2 vector bundle on $\mathbb{P}_3$, unique up to isomorphism. Curiously, there a way to globalise this construction over $\mathbb{P}_5$. In this part we will construct an SL_2 -equivariant rank 2 reflexive sheaf on $\mathbb{P}_5$ using classical techniques developped in mathematical instanton

studies. (Cf [Ba], [Tj]) Still consider vector spaces L and V of respective dimension 2 and 6. This construction will be essentially unique up to the SL_6 action. Indeed, it could be constructed from a tangent hyperplane to $G_\omega(3, 6)$ or like this:

Let's recall from [M-M] that $S_2L \otimes \bigwedge^2 V$ have an $SL_2 \times SL_6$ orbit made of net of alternating forms of constant rank 4. So let's $\beta \in S_2L \otimes \bigwedge^2 V$ be the element of this orbit. This element was considered in [M-dP] to construct a $\mathbb{P}_{11}$ containing their congruence, for instance, take the following:

$$\begin{pmatrix} 0 & u^2 & 2uv & v^2 & 0 & 0 \\ -u^2 & 0 & 0 & 0 & 0 & 0 \\ -2uv & 0 & 0 & 0 & 0 & u^2 \\ -v^2 & 0 & 0 & 0 & 0 & 2uv \\ 0 & 0 & 0 & 0 & 0 & v^2 \\ 0 & 0 & -u^2 & -2uv & -v^2 & 0 \end{pmatrix}$$

We can remark that β viewed as an element of $\bigwedge^2(L \otimes V)$ has rank 6 because it can be represented by:

$$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}$$

Denote by W the six dimensional image of this map, we will identify W with its dual via the induced alternating form. The inclusion $W \subset L \otimes V$ gives a map β' from W to $L \otimes \mathcal{O}_{\mathbb{P}_5}(1)$, and we can construct a complex:

$$L \otimes \mathcal{O}_{\mathbb{P}_5}(-1) \xrightarrow{t\beta'} W \otimes \mathcal{O}_{\mathbb{P}_5} \xrightarrow{\beta'} L \otimes \mathcal{O}_{\mathbb{P}_5}(1) \quad (6)$$

exact on the left, with middle cohomology a rank 2 reflexive sheaf K , and with right cohomology a sheaf $\mathcal{L}$ supported on a smooth rational cubic curve C isomorphic to $\mathcal{O}_{\mathbb{P}_1}(4)$. So we can compute from the complex (6) some of its invariants:

Corollary 4.7 *The sheaf K has rank 2, $c_1K = 0$, $c_2K = 2$, $c_3K = 0$, $c_4K = -15$. Its singular locus is the cubic curve C , and we have $H^0K(1) = 0$ and $H^0K(2) = 13$, and $H^1K = 1$.*

we have now a way to structurate the variation of the focal locus with respect to the choice of the quadric in th8 of [M-dP]. They are the vanishing locus of sections of the same bundle: $K(2)$.

References

- [Ba] Barth, W: Irreducibility of the space of mathematical instanton bundles with Rank 2, $c_2 = 4$. Math.Ann **258**, 81-106 (1981)
- [G-P] L.Gruson - C.Peskin: Courbes de l'espace projectif, variétés de sécantes. Enumerative geometry and classical algebraic geometry. Nice (1981) Prog Math **24** Birkhäuser.Boston 1982
- [I] A. Iliev: The SP_3 -grassmannian and duality for prime Fano threefolds of genus 9. Manuscripta math. **112**, 29-53 (2003)
- [I-R] A. Iliev - K. Ranestad: Geometry of the Lagrangian Grassmannian $LG(3,6)$ with applications to Brill-Noether loci. Mich. Math. Journal **53**, 383-417 (2005)
- [K] A. Kuznetsov: Hyperplane sections and derived categories. Izvestiya Mathematics **70**:3, 447-447 (2006)
- [M] L.Manivel: Configuration of lines and models of Lie algebras. Journal of Algebra **304**, 457-486 (2006)
- [M-M] L. Manivel - E. Mezzeti: On linear spaces of skew-symmetric matrices of constant rank. Manuscripta math. **117**, 319-331 (2005)
- [M-dP] Mezzeti - de Poi: Congruences of lines in $\mathbb{P}_5$, quadratic normality, and completely exceptional Monge-Ampère equations. Geom Dedicata **131**, 213-230 (2008)
- [Tj] Tjurin, A.N: On the superpositions of mathematical instantons. In Artin, Tate, J.(eds) Arithmetic and geometry. Prog.Math **36**, 433-450 (1983) . Birkhäuser.
- [W] J. Weyman: Cohomology of vector bundles and syzygies. **149** (2003) Cambridge Tracts in Mathematics.